

causal organism of little leaf of brinjal Full Version

Introduction to Little Leaf Disease of Brinjal

The causal organism of little leaf of brinjal (*Solanum melongena*) is a significant pathogen responsible for a devastating disease that impacts the growth, yield, and overall health of the plant. This disease is characterized by stunted plant growth, reduced leaf size, and distorted or misshapen foliage, ultimately leading to poor fruit development and significant economic losses for farmers. Understanding the causal organism behind this disease is crucial for effective management and control strategies to safeguard brinjal crops.

Overview of Little Leaf Disease

Symptoms of Little Leaf Disease

- Reduced leaf size, often less than half the normal size
- Yellowing or chlorosis of the leaves
- Stunted plant growth with shortened internodes
- Leaves may be curled or distorted
- Poor flowering and fruiting
- In severe cases, plants may die prematurely

Impact on Crop Production

Little leaf disease severely hampers the plant's ability to perform photosynthesis effectively, leading to reduced vigor and yield. The disease not only affects the individual plant but can spread rapidly within a crop, causing widespread damage. For farmers relying on brinjal as a staple or income-generating crop, controlling this disease is vital for economic sustainability.

The Causal Organism of Little Leaf of Brinjal

Identification of the Pathogen

The primary causal agent of little leaf disease in brinjal is a phytoplasma, a type of wall-less, obligate parasitic bacterium that infects plant phloem tissue. These microorganisms are not true bacteria but are classified as mollicutes, sharing characteristics with both bacteria and viruses. They are

transmitted by specific insect vectors, primarily leafhoppers, which feed on infected plants and subsequently spread the pathogen to healthy plants.

Phytoplasma: The Causal Organism

- **Taxonomy:** Belong to the class Mollicutes, order Acholeplasmatales, family 'Candidatus Phytoplasmataceae'.
- **Morphology:** Very small, pleomorphic, and wall-less organisms measuring approximately 200-800 nm in size.
- **Genetic Material:** Contain circular, single-stranded DNA genomes.
- **Lifestyle:** Obligate parasites residing in the phloem sieve elements of plants and the guts of insect vectors.

Role of Phytoplasma in Disease Development

The phytoplasma infects the phloem tissue of brinjal, disrupting normal nutrient translocation and hormonal balance. This interference results in abnormal growth patterns such as reduced leaf size and stunted growth. The pathogen's presence also triggers physiological changes, including the production of phytoplasma-specific symptoms like little leaves, witches' broom, and phyllody. The disease severity depends on factors such as the phytoplasma strain, plant age, environmental conditions, and vector population density.

Transmission of the Causal Organism

Vector Transmission

The primary mode of phytoplasma spread is through insect vectors, predominantly leafhoppers of the family Cicadellidae. These insects acquire the phytoplasma while feeding on infected plants and transmit it to healthy plants during subsequent feedings. The process involves:

- **Insect feeding on infected plant:** The leafhopper ingests the phytoplasma in the phloem sap.
- **Latency period:** The phytoplasma multiplies within the insect for a period before becoming transmissible.
- **Feeding on healthy plants:** The insect injects the phytoplasma into the phloem tissue of new plants, initiating infection.

Other Modes of Transmission

- **Vegetative propagation:** Use of infected cuttings or seedlings can spread the pathogen.
- **Mechanical transmission:** Rare, but possible through contaminated tools or handling.

Detection and Diagnosis of Little Leaf Disease

Field Diagnosis

Field symptoms such as small, chlorotic leaves, stunting, and distortion are initial indicators of infection. However, these symptoms can be confused with other diseases or nutrient deficiencies, necessitating laboratory confirmation.

Laboratory Techniques

- **Serological tests:** Enzyme-linked immunosorbent assay (ELISA) for detecting phytoplasma antigens.
- **DNA-based methods:** Polymerase chain reaction (PCR) using specific primers to identify phytoplasma DNA.
- **Electron microscopy:** Visualization of phytoplasma particles within phloem tissues.

Management and Control Strategies

Integrated Disease Management

- **Use of resistant varieties:** Breeding and selecting brinjal varieties less susceptible to phytoplasma infection.
- **Control of insect vectors:** Employing insecticides, biological control agents, and cultural practices to reduce leafhopper populations.
- **Sanitation:** Removing and destroying infected plants to prevent disease spread.
- **Crop rotation and field hygiene:** Avoiding planting brinjal in infested fields and practicing proper field sanitation.
- **Use of healthy planting material:** Ensuring seedlings are free from infection through certified nurseries.

Biological and Chemical Control

While chemical control of phytoplasma is challenging, targeting the insect vectors remains effective. Use of insecticides at appropriate timings can significantly reduce vector populations. Biological control agents like parasitic wasps and entomopathogenic fungi can also help manage leafhopper populations.

Conclusion

The causal organism of little leaf of brinjal is a phytoplasma, a wall-less, obligate parasitic microorganism that infects the plant's phloem tissue and causes characteristic symptoms such as reduced leaf size and stunted growth. Its transmission primarily occurs through leafhopper vectors, emphasizing the importance of integrated pest and disease management strategies. Accurate diagnosis using modern laboratory techniques is essential for effective control. Preventive measures, including resistant varieties, vector control, and sanitation, are vital to managing and mitigating the impact of this disease. Understanding the biology and transmission of the phytoplasma is crucial for developing sustainable solutions to protect brinjal crops from little leaf disease, ensuring better yields and economic stability for farmers.

Causal Organism of Little Leaf of Brinjal: An In-Depth Analysis

The health and productivity of brinjal (*Solanum melongena*), commonly known as eggplant, are vital to farmers and consumers worldwide, especially in tropical and subtropical regions. One of the most insidious threats to brinjal cultivation is the disease known as "little leaf," characterized by stunted growth, small and malformed leaves, and reduced fruit yield. At the heart of this disorder lies a specific causal organism that triggers these symptoms, and understanding this pathogen is crucial for effective management and control strategies. This article delves into the causal organism behind little leaf of brinjal, exploring its biology, mode of action, and implications for disease management.

Understanding Little Leaf Disease of Brinjal

Little leaf disease manifests primarily through morphological changes in the plant's foliage and growth pattern. Infected plants often exhibit:

- Reduced leaf size, often less than half the normal size
- Distorted leaf shape with thickened or curlings
- Stunted overall plant growth
- Reduced flowering and fruiting
- Chlorosis or yellowing in younger leaves

These symptoms translate into significant yield losses, making early diagnosis and understanding of the underlying cause essential for farmers and plant pathologists.

The Causal Organism: A Phytoplasma

What Are Phytoplasmas?

The primary causal agent of little leaf disease in brinjal is a specialized type of microorganism known as a phytoplasma. Phytoplasmas are tiny, wall-less bacteria classified within the class Mollicutes. Unlike typical bacteria, they lack a cell wall, making them pleomorphic (capable of changing shape), and they are obligate parasites, meaning they can only survive and multiply within the phloem tissue of host plants and insects.

These organisms are known to cause a variety of plant diseases worldwide, often characterized by symptoms such as leaf curl, virescence, phyllody, and stunting. In the case of brinjal, phytoplasma infection leads to the classic little leaf symptomatology.

Biology and Characteristics of Phytoplasmas

- **Size and Structure:** Phytoplasmas are extremely small, measuring approximately 200-800 nanometers in diameter, invisible under standard light microscopy. They are pleomorphic and lack a cell wall, which makes them resistant to some antibiotics and difficult to culture in laboratory media.
- **Genetic Makeup:** They possess a minimal genome, with a reduced set of genes necessary for parasitism and survival within the host. Molecular techniques, such as PCR and DNA sequencing, are often employed to identify specific phytoplasma strains.
- **Habitat:** They inhabit the phloem tissue of infected plants and are transmitted by certain insect vectors, predominantly leafhoppers, planthoppers, and psyllids.

Transmission of the Phytoplasma

The spread of little leaf disease is primarily mediated through insect vectors. The insect-plant interaction plays a pivotal role in the epidemiology of the disease:

- **Insect Vectors:** Several species of leafhoppers (family Cicadellidae), particularly the *Empoasca* spp., are known vectors. These insects acquire the phytoplasma when feeding on infected plants and subsequently transmit it to healthy plants during subsequent feeding.
- **Mechanism of Transmission:** The phytoplasma resides in the insect's salivary glands. When the insect feeds on a healthy brinjal plant, the pathogen is inoculated into the plant's phloem tissue.

- Other Modes of Spread: While insect transmission is predominant, the disease can also spread through infected planting material, contaminated tools, or via grafting practices if infected tissue is used.

Symptoms Attributed to Phytoplasma Infection

The infection of brinjal plants with phytoplasmas manifests in distinctive symptoms that aid in diagnosis:

- Little Leaf: The most characteristic symptom. The leaves are abnormally small, often less than 2-3 inches in length, and may be thickened or leathery.
- Leaf Curl and Virescence: Some infected plants show curling of leaf margins and light green to yellowish coloration.
- Stunting: The overall plant growth is suppressed, resulting in shorter stature and fewer branches.
- Flower and Fruit Abnormalities: Infected plants may produce fewer flowers, and fruits can be deformed or reduced in size.
- Vascular Discoloration: In advanced stages, the phloem tissue may show discoloration due to the pathogen's proliferation.

Diagnosis and Identification

Accurate diagnosis of phytoplasma-induced little leaf is crucial for implementing control measures. Traditional visual assessment can be misleading; thus, laboratory techniques are employed:

- Serological Tests: ELISA (Enzyme-Linked Immunosorbent Assay) can detect phytoplasma-specific antigens.
- Molecular Techniques: PCR (Polymerase Chain Reaction) using phytoplasma-specific primers provides definitive identification. Quantitative PCR (qPCR) can estimate pathogen load.
- Electron Microscopy: While not routine, it can visualize phytoplasma filaments within the phloem tissue.

Implications for Disease Management

Understanding that phytoplasmas are the causal agents of little leaf disease informs several management strategies:

- Use of Disease-Free Planting Material: Since infected tissue can harbor the pathogen, sourcing

healthy seedlings is essential.

- Control of Insect Vectors: Managing leafhopper populations through insecticides, traps, or cultural practices reduces transmission risk.
- Cultural Practices: Removing infected plants, crop rotation, and managing weeds that may host vectors or phytoplasma.
- Chemical Control: No effective cure exists for phytoplasma infection; however, controlling vectors can limit spread.
- Breeding for Resistance: Developing and deploying brinjal varieties resistant or tolerant to phytoplasma infection is an ongoing research focus.

Research and Future Perspectives

Advances in molecular biology have enhanced understanding of phytoplasma diversity, epidemiology, and interactions with hosts. Current research aims to:

- Identify resistant brinjal cultivars.
- Develop biological control agents targeting insect vectors.
- Enhance rapid, field-level diagnostic tools.
- Explore genetic engineering approaches to confer resistance.

Furthermore, integrated disease management strategies combining cultural, biological, and chemical methods offer the best prospects for controlling little leaf disease effectively.

Conclusion

The causal organism of little leaf of brinjal is a phytoplasma—a wall-less, obligate parasitic bacterium that resides within the plant's phloem tissue. Transmitted chiefly by leafhoppers, these microorganisms induce characteristic symptoms such as reduced leaf size, plant stunting, and poor fruiting. Recognizing the role of phytoplasmas in this disease underscores the importance of integrated management practices, including vector control, use of healthy planting material, and ongoing research into resistant varieties. As our understanding deepens, the prospects for mitigating the impact of little leaf disease on brinjal cultivation become increasingly promising, ensuring better yields and sustainability for farmers worldwide.

Question What is the causal organism responsible for little leaf disease in brinjal? The causal organism of little leaf disease in brinjal is the phytoplasma, specifically the aster yellows phytoplasma group. **How does the phytoplasma cause little leaf symptoms in brinjal plants?** The phytoplasma infects the phloem tissue, disrupting nutrient transport and causing stunted growth, small leaves, and reduced plant vigor characteristic of little leaf disease. **What are the primary symptoms of little leaf disease in brinjal?** Symptoms include small, distorted, and pale green leaves,

stunted plant growth, and reduced fruiting, often leading to significant yield loss. Which insects are known vectors transmitting the phytoplasma causing little leaf in brinjal? Leafhoppers, particularly species like *Empoasca* spp., are primary vectors transmitting the phytoplasma to brinjal plants. How can farmers manage or control the spread of little leaf disease in brinjal? Management includes controlling leafhopper populations through insecticides, removing infected plants, using resistant varieties, and practicing crop rotation to reduce vector populations. Are there resistant varieties of brinjal to little leaf disease? Yes, some brinjal varieties exhibit resistance or tolerance to the disease, but identification and adoption depend on local conditions and availability. Is little leaf disease in brinjal a viral, bacterial, or fungal disease? Little leaf in brinjal is caused by a phytoplasma, which is a type of wall-less bacteria-like organism, not a virus or fungus. Can chemical control cure little leaf disease in brinjal? Chemical control is ineffective against phytoplasmas directly; management focuses on controlling vectors and cultural practices to reduce disease incidence. What is the importance of early detection of little leaf in brinjal for effective management? Early detection allows timely removal of infected plants and vector control measures, helping to prevent the spread and minimize crop losses.

Related keywords: Brinjal, Little Leaf Disease, Causal Organism, Phytoplasma, Aster Yellows, Mycoplasma-like organisms, Eggplant, Plant Pathology, Disease Transmission, Insect Vector

DICHOTOMOUS KEY REGENTS QUESTION

dichotomous key regents question ? a phrase that resonates deeply with students preparing for science exams, especially in biology. These questions are fundamental in understanding how scientists classify and identify organisms systematically. Mastering dichotomous keys is essential for success in regents exams, as they test your ability to analyze, differentiate, and logically deduce the identity of various organisms based on observable traits. This comprehensive guide aims to equip students with an in-depth understanding of dichotomous key regents questions, strategies for answering them effectively, and tips to excel in your assessments.

Understanding the Concept of a Dichotomous Key

What Is a Dichotomous Key?

A dichotomous key is a tool used by biologists and students to identify unknown organisms or objects through a series of choices that lead to the correct identification. The term "dichotomous" means "divided into two parts," reflecting the structure of the key, which presents two contrasting options at each step.

Key features of a dichotomous key include:

- A sequence of paired statements or questions
- Each choice narrows down the possibilities
- The process continues until the organism is identified accurately
- Often used for plants, animals, insects, and other biological specimens

How Does a Dichotomous Key Work?

The process involves:

- Observing the organism's features
- Selecting the statement that best describes the organism's current trait
- Moving to the next set of choices based on that selection
- Repeating until reaching the final identification

Example flow:

- Does the organism have leaves? ? Yes ? Proceed to step 2
- Are the leaves needle-like? ? Yes ? It's a pine tree

OR

- Does the organism have leaves? ? No ? Proceed to step 3

This step-by-step approach simplifies complex identification tasks into manageable decisions.

Common Types of Regents Questions on Dichotomous Keys

Multiple-Choice Questions

These questions present a scenario where you choose the correct next step or identification based on a description or image.

Example:

Using a dichotomous key, you observe an insect with wings and six legs. The key asks:

- a) Does the insect have a pair of antennae? ? Yes
- b) Does the insect have a segmented body? ? Yes

What is the insect?

Options:

- A) Beetle
- B) Butterfly
- C) Ant
- D) Fly

Answer: D) Fly

Scenario-Based Questions

These involve analyzing a description or picture, then selecting the correct sequence of choices in the key to identify the organism.

Example:

You are shown an unknown plant. The question asks:

- Step 1: Does the plant have flowers? ? Yes
- Step 2: Are the flowers grouped in clusters? ? No
- Step 3: Are the leaves broad or narrow? ? Broad

What is the plant?

Diagram or Image Interpretation Questions

Questions may include diagrams of organisms or parts, requiring you to interpret features and follow the key accordingly.

Strategies for Answering Dichotomous Key Regents Questions Effectively

1. Carefully Read Each Step

- Pay attention to the specific traits described in each choice.
- Look for keywords that match the organism's features.

2. Observe the Organism or Image Accurately

- Note details such as shape, size, color, number of legs, presence of wings, leaf arrangement, etc.
- Use a systematic approach: examine one feature at a time.

3. Use Process of Elimination

- Narrow down options by eliminating choices that do not match the observed traits.
- Remember, each step should refine your options further.

4. Follow the Logical Flow

- Move sequentially through the key based on your observations.
- Do not skip steps or make assumptions beyond what the traits suggest.

5. Practice with Sample Questions

- Familiarize yourself with common organisms and features.
- Practice with past regents or sample dichotomous keys to build confidence.

Common Mistakes to Avoid in Regents Questions on Dichotomous Keys

- **Misreading Traits:** Always double-check the organism's features before making a choice.
- **Ignoring Key Words:** Words like "needle-like," "segmented," "broad," or "compound" are vital.
- **Skipping Steps:** Follow each step carefully; missing a choice can lead to incorrect identification.
- **Assuming Traits Not Observed:** Base your decisions solely on observable features, not assumptions.

Sample Practice Question and Step-by-Step Solution

Question:

Using a dichotomous key, identify this organism: It has a segmented body, six legs, and two antennae. The organism has wings, and the wings are covered with scales. Based on the key, which organism is this?

Step-by-step solution:

- Does the organism have wings?
- Yes ? Proceed to next step
- Are the wings covered with scales?
- Yes ? The organism is a butterfly or moth
- Does the organism have a segmented body with six legs?
- Yes ? It is likely a butterfly

Final identification: Butterfly

Preparing for Regents Exams: Tips and Resources

Study Tips

- Review common organisms and their traits.
- Practice identifying features in diagrams and photographs.
- Create your own dichotomous keys for familiar objects or animals.
- Use flashcards to memorize key traits associated with different species.

Useful Resources

- Past regents exams with dichotomous key questions
- Biology textbooks and lab manuals
- Online quizzes and interactive dichotomous key tools
- Educational videos explaining how to use and construct dichotomous keys

Conclusion

Mastering dichotomous key regents questions is vital for success in biology exams. By understanding how dichotomous keys function, practicing systematically, and paying close attention to observable traits, students can improve their identification skills and confidence. Remember, the key to excelling is patience, careful observation, and consistent practice. Use this guide as a resource to sharpen your skills and approach each regents question with clarity and precision. Good luck!

Understanding and Mastering Dichotomous Key Regents Questions: A Comprehensive Guide

When preparing for standardized exams like the Regents, students often encounter a variety of question types designed to assess their understanding of biological classification and identification. Among these, dichotomous key regents questions are particularly common and can be challenging if not approached strategically. These questions test your ability to interpret and apply dichotomous keys?tools used by biologists to identify organisms based on a series of paired choices. Mastering these questions requires a clear understanding of how dichotomous keys function, as well as effective strategies for navigating them during exams.

What Is a Dichotomous Key?

A dichotomous key is a tool that allows for the identification of organisms or objects through a sequence of choices. Each step presents two contrasting statements (or ?couplets?) about a characteristic of the organism. Based on your answer, you move to the next set of choices until you reach the final identification.

Key features of a dichotomous key include:

- Paired statements: Each step offers two options that are mutually exclusive.
- Sequential decision-making: Choices lead you progressively toward the correct identification.
- Hierarchical structure: The key narrows down possibilities at each step, making the process systematic.

Why Are Dichotomous Key Questions Important in the Regents Exam?

Dichotomous key questions are integral because they assess a student's ability to:

- Analyze physical features or traits of organisms.
- Apply logical reasoning to differentiate between similar species or objects.
- Follow complex instructions to arrive at a correct identification.

Successfully answering these questions demonstrates not only memorization but also critical thinking and process skills that are vital in biological sciences.

Breaking Down the Structure of a Dichotomous Key Regents Question

A typical dichotomous key question on the Regents exam may present an image, description, or list of characteristics, and then ask you to:

- Choose the correct next step in the key.
- Identify an organism based on a series of choices.
- Determine which statement correctly differentiates two similar species.

Understanding how these questions are structured helps you approach them methodically.

Step-by-Step Approach to Answering Dichotomous Key Questions

- Carefully Read the Question and All Provided Information
- Look at images, descriptions, or diagrams.
- Note specific traits highlighted, such as leaf shape, color, number of legs, or habitat.
- Identify the Relevant Characteristics
- Focus on the features that are explicitly mentioned or depicted.
- Ignore irrelevant information to avoid confusion.
- Follow the Logical Sequence of the Key

- Recognize that each choice in the key leads to a different subsequent step.
- Use process of elimination to narrow down options.

- Match Characteristics to the Correct Statement

- Determine which of the paired statements accurately describes the organism based on the traits provided.
- Confirm that your choice aligns with the organism's features.

- Use the Process of Elimination

- Eliminate options that do not fit the traits.
- Narrow your choices systematically until you reach the final identification.

- Double-Check Your Answer

- Review the characteristics and ensure consistency with your selected choice.
- Verify that the path through the key makes logical sense.

Common Challenges and How to Overcome Them

Challenge 1: Confusing Similar Traits

- Solution: Pay close attention to specific details, such as the number of petals, leaf arrangement, or presence of certain structures.

Challenge 2: Misreading Paired Statements

- Solution: Read each statement carefully, noting keywords that differentiate the options.

Challenge 3: Forgetting the Sequence

- Solution: Practice with sample keys to familiarize yourself with the flow and structure of decision trees.

Practice Tips for Mastering Dichotomous Key Questions

- Use Visual Aids: Practice with diagrams and actual specimens if possible.
- Create Your Own Keys: Try designing simple dichotomous keys for familiar objects to understand the decision process.
- Work Through Past Regents Questions: Review previous exams to recognize patterns and question formats.
- Time Management: Practice under timed conditions to improve efficiency during the exam.
- Discuss with Peers or Teachers: Collaborative learning can clarify confusing traits and decision points.

Example Walkthrough of a Typical Dichotomous Key Question

Suppose a question presents an image of a leaf and asks:

Based on the leaf's shape and vein pattern, which of the following organism identifications is correct?

Possible choices are:

- A) Maple tree
- B) Oak tree
- C) Elm tree
- D) Birch tree

Approach:

- Identify traits: Is the leaf lobed? Are the veins branching?
- Cross-reference these traits with the key choices:
 - Maples have lobed leaves with palmate veins.
 - Oaks have lobed leaves but with different vein patterns.
 - Elms have serrated edges and pinnate veins.
 - Birches have simple, pointed leaves with specific vein arrangements.

- Match traits: If the leaf has a palmate vein pattern with lobed margins, the answer is A) Maple tree.

Final Tips for Success on Dichotomous Key Questions

- Stay Calm and Focused: Carefully read each step and avoid rushing.
- Practice Regularly: Familiarity reduces errors and increases confidence.
- Understand the Traits: Know the key features that distinguish major groups of organisms.
- Use Process of Elimination: When unsure, eliminate clearly incorrect options to narrow your choices.

Conclusion

Dichotomous key regents questions are designed to challenge your ability to analyze biological traits systematically. By understanding how dichotomous keys work, developing a logical approach to navigating them, and practicing extensively, you can improve your accuracy and confidence. Remember, these questions are less about memorization and more about applying your observational and reasoning skills. With diligent preparation and strategic thinking, you'll be well-equipped to master these questions and excel on your Regents exam.

Happy studying, and good luck on your biology journey!

Question What is a dichotomous key and how is it used in regents exams? A dichotomous key is a tool used to identify organisms or objects by answering a series of paired questions that lead to the correct identification. In regents exams, students often interpret or create dichotomous keys to classify biological specimens. How can I effectively approach a regents question involving a dichotomous key? Read each question carefully, follow the logical flow of the key, eliminate options based on the characteristics described, and choose the identification that matches the organism or object described in the question. What are common mistakes to avoid when answering dichotomous key questions on the regents? Common mistakes include misreading characteristics, skipping steps, selecting the wrong trait at a decision point, and not carefully comparing the organism's features to the key options. Can you provide an example of a typical regents question involving a dichotomous key? Sure! For example: 'Using the dichotomous key, identify the organism based on these traits: has wings, and is a herbivore. Which organism is it?' The key would guide you through choices to arrive at the correct organism. Why are dichotomous keys important in biology regents exams? They help students develop skills in classification, observation, and logical thinking, which are essential for understanding biological diversity and proper identification of species. What should I review before taking a regents exam with dichotomous key questions? Review the structure and terminology of dichotomous keys, practice with sample keys, and familiarize yourself with common organism characteristics used in

classification. Are there tips for memorizing traits used in dichotomous keys for regents exams? Yes, creating flashcards for key traits, practicing with sample keys regularly, and understanding the significance of each characteristic can improve recall and application during the exam. How can I practice improving my skills with dichotomous keys for regents preparation? Use past exam questions, practice with biological specimens or diagrams, create your own keys for different organisms, and seek feedback from teachers or study groups to reinforce understanding.

Related keywords: dichotomous key, regents exam, biology, identification, taxonomy, classification, plant identification, animal identification, science assessment, educational resources

Read the full article online: [DICHOTOMOUS KEY REGENTS QUESTION](#)